

Submillimeter Array Observations of Disks in the SR 24 Multiple Star System¹

Sean M. Andrews & Jonathan P. Williams

Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, HI 96822

andrews@ifa.hawaii.edu, jpw@ifa.hawaii.edu

ABSTRACT

We present high-resolution aperture synthesis images from the Submillimeter Array of the 225 GHz (1.3 mm) continuum and ^{12}CO $J = 2 - 1$ line emission from the disks around components of the hierarchical triple system SR 24, located in the Ophiuchus star-forming region. The most widely separated component, SR 24 S (with a projected semimajor axis $a = 832$ AU), has a circumstellar disk with properties typical of those around single T Tauri stars. The binary SR 24 N ($a = 32$ AU) is undetected in the continuum, but has strong, resolved CO emission which likely originates in a circumbinary disk with a central gap. The data constrain the total disk mass in the SR 24 N system to be $\lesssim 10^{-3} M_{\odot}$ and indicate that the depletion of CO onto dust grains is not more than 100 times larger than the mean value in the interstellar medium. The SR 24 N disk is unusual in that it is only detected in line emission. It is possible that other low mass disks around binaries and single stars may have been missed in single-dish continuum surveys.

Subject headings: binaries: close — stars: individual (SR 24) — stars: circumstellar matter

1. Introduction

Much progress has been made in constraining the basic physical properties of circumstellar disks with the inclusion of observations at (sub)millimeter wavelengths, beginning with a set of large 1.3 mm continuum surveys (Beckwith et al. 1990; André & Montmerle 1994; Osterloh & Beckwith 1995). At these long wavelengths, the thermal continuum emission from the dust disk is assumed to be optically-thin and thus provides a direct constraint on the total particle mass (Beckwith et al. 1986). Moreover, the (sub)millimeter spectrum is rich with rotational transitions of various molecules, presenting opportunities to study kinematics and chemistry (e.g., Qi et al.

¹The Submillimeter Array is a joint project between the Smithsonian Astrophysical Observatory and the Academia Sinica Institute of Astronomy and Astrophysics, and is funded by the Smithsonian Institution and the Academia Sinica.

2003). Because an interferometer is required to disentangle the line emission from the disk and molecular cloud, only a handful of disks have constraints on the molecular gas phase.

The metamorphosis of a circumstellar disk into a planetary system can be inhibited if the local environment is hostile enough to severely disturb or destroy the disk. A perhaps common example is photoevaporation in the vicinity of a massive star (e.g., Bally et al. 1998). Another is dynamical disruption due to another star, either in a multiple system or a dense cluster environment. The frequency of multiple systems among young stars is at least equal to that of main sequence stars in the field, and possibly significantly greater (Mathieu 1994; Mathieu et al. 2000). Numerical simulations demonstrate that the stability of circumstellar material in multiple star systems is jeopardized (Papaloizou & Pringle 1977; Lin & Papaloizou 1993; Artymowicz & Lubow 1994). These studies show that, despite the dynamical interactions between disks and stars, individual *circumstellar* disks can survive, albeit in truncated form, and large gaps would be produced in the *circumbinary* disks. Single-dish continuum searches have confirmed that close binaries ($a \lesssim 50$ AU) have significantly lower (sub)millimeter luminosities than wider binaries or single stars (Osterloh & Beckwith 1995; Jensen, Mathieu, & Fuller 1994). However, the luminosities are still consistent with dynamically-carved gaps in circumbinary disks, and not necessarily the absence of disks altogether (Jensen, Mathieu, & Fuller 1996). Many young binaries have clear signatures of inner disks around at least one of the components (Jensen & Mathieu 1997). Recent evidence shows that widely-separated components are able to retain significant disks (Jensen & Akeson 2003).

The focus of this Letter is the triple system SR 24 (HBC 262, Haro 1–7), located in the Ophiuchus star-forming region ($d = 160$ pc; Chini 1981). For convenience we refer to the primary as SR 24 S and the binary as SR 24 N. The projected separation of SR 24 N/S is $5''.2$ at $PA = 348^\circ$ (Reipurth & Zinnecker 1993). SR 24 N itself is a binary with a projected separation of $0''.197$ at $PA = 87^\circ$ (Simon et al. 1995). Both SR 24 N and S are known to be classical T Tauri stars with emission lines (Cohen & Kuhl 1979), Li absorption (Martín et al. 1998), and infrared excesses (Greene et al. 1994). We present new high-resolution observations of the SR 24 system in the 1.3 mm continuum and CO $J = 2 - 1$ line, and discuss their implications on understanding disk structure and planet formation in multiple star systems.

2. Observations and Data Reduction

Millimeter interferometric observations of the SR 24 system were obtained on 2004 August 2 and 21 with the Submillimeter Array (SMA; Ho, Moran, & Lo 2004) on Mauna Kea, Hawaii. The data were taken with 7 antennas and double-sideband receivers with $\nu_{\text{IF}} = 225.494$ GHz. The ^{12}CO $J = 2 - 1$ line at 230.538 GHz was centered in the upper sideband. The selected correlator setup gave 2 GHz of continuum bandwidth in each sideband and a spectral resolution of 1.06 km s^{-1} (~ 0.8 MHz) per channel. During the observations, the zenith opacity was ~ 0.12 and the median system temperature was ~ 160 K. The combined compact (August 2) and extended (August 21) array configurations resulted in projected baselines from 9 to 186 m. Observations

of SR 24 (with the phase center set at the position of SR 24 S: $\alpha=16^{\text{h}}26^{\text{m}}58^{\text{s}}.5$, $\delta=-24^{\circ}45'36''.67$, J2000) were interleaved with two quasars (J1743–038 and NRAO 530) for use in gain calibration. Additional observations of Jupiter and Uranus were obtained for bandpass and flux calibration. The total on-source integration times for SR 24 were 1.7 and 3.3 hours for the compact and extended configurations, respectively.

The data were edited and calibrated using the IDL-based MIR software package.² The bandpass response was determined from the Jupiter and Uranus observations. Complex gain calibration was conducted with both quasars, and the absolute flux calibration was set by J1743–038 ($F_{\nu} = 2.0$ Jy). The resulting quasar flux densities are in excellent agreement with several independent measurements conducted throughout 2004 August, from which we infer that the absolute flux calibration is accurate to $\sim 20\%$. The resulting rms phase noise for the quasars increased from $\sim 15^{\circ}$ to 60° from the shortest to the longest projected baselines. Standard imaging and deconvolution was conducted with the MIRIAD software package. All maps were obtained using natural *uv*-weighting.

3. Results

The millimeter continuum map of the SR 24 system has a FWHM resolution of $2''.3 \times 1''.2$ (beam PA = 43°) and rms noise level of 2 mJy beam^{-1} . A continuum source with a flux density of 68 mJy is detected at the position of SR 24 S. An elliptical gaussian fit to the source in a higher resolution map ($1''.4 \times 1''.0$, beam PA = 43°) made using only the extended array data shows that it is resolved, with a deconvolved FWHM size of $1''.23 \pm 0''.02 \times 0''.29 \pm 0''.01$ at PA = 28° . The SR 24 S flux density is a factor of 3 to 4 lower than previous single-dish estimates, where the FWHM beam sizes were $12 - 18''$ (Reipurth et al. 1993; André & Montmerle 1994; Nürnberger et al. 1998). The Fourier sampling of the SMA indicates that our sensitivity is decreased to only 10% of the peak at those single-dish beam sizes (e.g., Wilner & Welch 1994). Therefore, the single-dish observations likely suffer contamination from extended emission due to distributed dust, e.g., an envelope around the entire SR 24 system. No continuum emission is detected in the vicinity of SR 24 N to a 3σ upper limit of 6 mJy .

Channel maps of the CO line at the same FWHM resolution as the continuum data have an rms noise level of $0.13 \text{ Jy beam}^{-1}$ in each channel. There is weak CO emission coincident with SR 24 S which has a spatially-averaged peak brightness temperature $T_b = 3 \text{ K}$ at LSR velocity $v = 6 \text{ km s}^{-1}$. The velocity-integrated ($\Delta v = 2.1 \text{ km s}^{-1}$) line has an intensity $I_{\text{CO}} \approx 4 \text{ K km s}^{-1}$. A much stronger CO line is detected at the SR 24 N position, with a spatially-averaged peak $T_b = 5.7 \text{ K}$ at $v = 7 \text{ km s}^{-1}$ and a velocity-integrated intensity $I_{\text{CO}} = 16.8 \text{ K km s}^{-1}$. The line emission for SR 24 N is spatially resolved, and a fit with an elliptical gaussian gives a deconvolved FWHM size of

²<http://cfa-www.harvard.edu/~cqj/mircook.html>

$3''.2 \times 1''.9$ at $\text{PA} = 45^\circ$.

Figure 1 shows the millimeter continuum map in grayscale, with overlaid contours of integrated CO emission. The stellar positions have been determined from a *V*-band (F606W) image of SR 24 from the *Hubble Space Telescope* (HST) archive. While the positions of the stars relative to one another are known to very high accuracy, the absolute astrometric precision of the HST image is comparatively poor ($0''.7$; Baggett et al. 2002) due to the small field of view. The astrometric accuracy of the SMA data was found to be $\sim 0''.1$ by comparing the measured positions of the gain calibrators with their VLA coordinates. The line and continuum peaks are thus at the positions of SR 24 N and S within the uncertainties.

4. Discussion

The disk masses in the SR 24 system can be computed by assuming that the millimeter continuum emission originates in an optically-thin, isothermal region with the interstellar gas to dust mass ratio (~ 100). Using $T = 30$ K and an opacity of $0.02 \text{ cm}^2 \text{ g}^{-1}$ (Hildebrand 1983), the SR 24 S disk has $M_{\text{disk}} = 0.01 M_\odot$, which is roughly the mass obtained by augmenting the abundances of the planets in the solar system to cosmic values (the minimum mass solar nebula; e.g., Weidenschilling 1977). The mass and projected diameter (190 AU) of the SR 24 S disk are consistent with typical single T Tauri stars (e.g., Dutrey et al. 1996). With the same parameters, we can constrain the SR 24 N disk(s) to have $M_{\text{disk}} \leq 9.0 \times 10^{-4} M_\odot$ (3σ), which is roughly the mass of Jupiter.

However, the strong CO emission at the position of SR 24 N indicates that there is disk material around this binary. The velocity gradient across the emission is low ($\Delta v \leq 2 \text{ km s}^{-1}$), and therefore it is not likely to originate in a molecular outflow. There are no bright peaks similar to the emission coincident with SR 24 N in the primary beam ($2''.5$) of the line maps, making the probability of a chance alignment of cloud material negligible. There are slight position-velocity offsets in the CO emission which are consistent with Keplerian rotation centered on the binary, but data with higher spectral resolution would be required to address this quantitatively. In most disks, the low energy rotational lines of CO are expected to be optically-thick (Beckwith & Sargent 1993). Therefore, we can derive a lower limit on the H_2 gas mass by treating the line as if it were optically-thin, following the method of Scoville et al. (1986).³ For $T = 30$ K, the SR 24 N disk has a gas mass $M_g \geq 2.8 \times 10^{-9} X_{\text{CO}}^{-1} M_\odot$, where X_{CO} is the CO abundance relative to H_2 ($X_{\text{CO}} \equiv [\text{CO}/\text{H}_2]$). The upper limit from the dust and the lower limit from the gas imply $2.8 \times 10^{-9} X_{\text{CO}}^{-1} < M_{\text{disk}} (M_\odot) < 9.0 \times 10^{-4}$, which gives a 3σ lower limit on the CO abundance, $X_{\text{CO}} > 3.1 \times 10^{-6} = 3.1 \times 10^{-2} X_{\text{CO,ism}}$, where $X_{\text{CO,ism}} = 10^{-4}$. For temperatures between

³Here we also assume a 10% He fraction and that the excitation temperature of the gas and the thermal temperature of the dust grains are equal.

$T = 20 - 100$ K, the limit on X_{CO} varies from $0.02 - 0.26 X_{\text{CO,ism}}$, showing that the CO depletion is no more than 100 in the SR 24 N disk. In fact, these measurements are consistent with there being no depletion compared to the ISM in the SR 24 N disk. This contrasts with the SR 24 S disk where the ratio of the CO line intensity to the continuum flux density is more than a factor of 100 lower and the CO depletion is $\sim 10^3$ ($T = 30$ K), similar to other circumstellar disks.

The broadband spectral energy distributions (SEDs) of SR 24 N and S, shown in Figure 2, are similar shortward of $20\ \mu\text{m}$. The infrared data, taken from the work of Greene et al. (1994), indicate that both components are typical Class II sources. The infrared excesses out to $20\ \mu\text{m}$ coupled with the strong $\text{H}\alpha$ emission lines are clear indications that both sources harbor actively accreting disks out to radii of at least 1 AU. The semimajor axis of the SR 24 N binary is $a = 32$ AU. Dynamical simulations of binaries with disks show that the individual circumstellar disks will be rapidly truncated at a fraction ($\sim 0.2 - 0.4$, depending primarily on the orbital eccentricity) of the semimajor axis (Artymowicz & Lubow 1994). In the case of SR 24 N, the truncation radius is estimated to lie between 6 and 12 AU. Assuming radial temperature profiles of typical optically-thick Class II disks (e.g., Beckwith et al. 1990), the truncated SR 24 N circumstellar disks should be able to produce thermal excesses out to $\sim 100\ \mu\text{m}$, which is consistent with the SED.

The same dynamical process which truncates the circumstellar disks in the binary will open a large gap in the circumbinary disk to a radius a factor of $2 - 3\times$ the semimajor axis (Artymowicz & Lubow 1994), meaning $\sim 60 - 100$ AU for SR 24 N. The observed CO emission, which extends to a radius of ~ 250 AU, likely originates in the remnant outer portion of the SR 24 N circumbinary disk. Dynamical theory would predict that the SR 24 N disk structure consists of at least one circumstellar accretion disk with radius $\lesssim 12$ AU and a circumbinary ring which extends out from ~ 80 AU. The observations confirm the presence of an inner disk and a large gas disk extending to 250 AU, but cannot constrain a gap size if one indeed exists. Since the average density in a disk is expected to decrease with radius, the CO depletion may be less in the outer regions. If the CO emission around SR 24 N arises in an outer circumbinary ring, this may explain the very different relative strengths of the line and continuum emission in the SR 24 N and S disks. Because they are uncontaminated by emission from an inner disk, circumbinary structures like that around SR 24 N may be a good way to directly probe physical conditions in the outermost regions of disks.

Assuming the SR 24 N system has a reasonably eccentric orbit ($e \sim 0.4$), the upper limit on the continuum flux density is marginally consistent with the simple disk + gap models computed by Jensen, Mathieu, & Fuller (1996). Therefore, a gap in the circumbinary disk may be all that is required to account for the non-detection in the continuum. If the standard form of the mass surface density ($\Sigma \propto r^{-3/2}$) is unaffected by the clearing of the gap in the circumbinary disk, the fraction of the total mass removed is roughly a factor of 2 (independent of the normalization of Σ). This implies that the “undisturbed” disk structure around SR 24 N was not very massive; at most a few times Jupiter’s mass, and much less massive than the SR 24 S disk. Alternatively, the SR 24 N disk may have evolved faster than the SR 24 S disk.

A few other young multiple star systems have been observed with millimeter interferometers. At first glance, one of the most interesting is the UZ Tau quadruple system, observed by Jensen, Koerner, & Mathieu (1996). The widely-separated primaries SR 24 S and UZ Tau E⁴ both have roughly minimum mass solar nebula disks detected in the continuum and CO. However, the $a \sim 50$ AU binary UZ Tau W has weak, unresolved ($\lesssim 70$ AU) continuum emission and no CO line emission associated with it: essentially the complete opposite of SR 24 N. The small apparent size of the UZ Tau W continuum emission led Jensen, Koerner, & Mathieu (1996) to conclude that individual circumstellar disks contributed the continuum emission, whereas the gas around SR 24 N most likely arises in a circumbinary disk. At the other extreme is GG Tau A, which has an essentially identical semimajor axis ($a \sim 36$ AU) as SR 24 N and similar signatures of inner disk accretion, but a very massive ($M_d \approx 0.1 M_\odot$) circumbinary ring well detected in both line and continuum emission (Koerner, Sargent, & Beckwith 1993; Dutrey, Guilloteau, & Simon 1994). The disk around the UY Aur binary ($a \sim 120$ AU) is an intermediate case with extensive, bright CO emission and comparatively little continuum emission (Duvert et al. 1998). Relatively massive circumbinary disks are also found around spectroscopic binaries (e.g., GW Ori; Mathieu et al. 1995) and other triple systems (e.g., T Tau; Weintraub, Masson, & Zuckerman 1989).

There is obviously a range of disk structures possible around multiple star systems. With so few objects observed at high resolution, the cause of this diversity remains unclear. The differences could be due to the evolutionary state of the system, or simply the (presumed) variety in the orbital parameters of the stellar components. The strong CO detection of the SR 24 N disk in the absence of continuum emission introduces the intriguing possibility that other low mass disks, around both single and multiple stars, may have been missed in even the most sensitive single-dish continuum searches to date. In spite of that possibility, the low disk masses inferred from (sub)millimeter observations around most binaries with separations similar to SR 24 N suggest that planet formation would not be possible given our current theoretical understanding. On the contrary, the more widely-separated components in multiple star systems, like SR 24 S, appear to harbor disks which are just as capable of forming planetary systems as single stars (see also Jensen, Mathieu, & Fuller 1996; Jensen & Akeson 2003).

5. Summary

We have presented high-resolution aperture synthesis images from the SMA interferometer of the disks in the SR 24 triple system in Ophiuchus. The comparatively isolated component in the system, SR 24 S, harbors a typical circumstellar disk. On the other hand, the binary SR 24 N is only detected in CO line emission and not the continuum. We suggest that SR 24 N is surrounded by at least one circumstellar disk as well as a circumbinary gas disk, presumably with

⁴This star is actually a spectroscopic binary (Mathieu, Martín, & Magazzù 1996), and thus is another example of a circumbinary disk.

a dynamically-carved gap. Little is known about the molecular gas phase in disks around single stars, let alone high-order multiple systems. Because multiple star systems like SR 24 are coeval, the time evolution factor is removed, so that the variety in the observed disk structures is likely due to the effects of the local environment.

We are grateful for the assistance and expertise of the SMA staff, and in particular for the advice of Alison Peck, David Wilner, Shigehisa Takakuwa, and Chunhua Qi. We also thank an anonymous referee and Mike Liu for useful suggestions which improved this paper. This work was supported by NSF grant AST-0324328.

REFERENCES

- André, P., & Montmerle, T. 1994, *ApJ*, 420, 837
- Artymowicz, P., & Lubow, S. H. 1994, *ApJ*, 421, 651
- Baggett, S. et al. 2002, *HST* WFPC2 Data Handbook, Version 4.0, ed. B. Mobasher (Baltimore: STScI)
- Bally, J., Testi, L., Sargent, A., & Carlstrom, J. 1998, *AJ*, 116, 854
- Beckwith, S. V. W., Sargent, A. I., Scoville, N. Z., Masson, C. R., Zuckerman, B., & Phillips, T. G. 1986, *ApJ*, 309, 755
- Beckwith, S. V. W., Sargent, A. I., Chini, R. S., & Güsten, R. 1990, *AJ*, 99, 924
- Beckwith, S. V. W., & Sargent, A. I. 1993, *ApJ*, 402, 280
- Chini, R. 1981, *A&A*, 99, 346
- Cohen, M., & Kuhl, L. V. 1979, *ApJS*, 41, 743
- Dutrey, A., Guilloteau, S., & Simon, M. 1994, *A&A*, 286, 149
- Dutrey, A., Guilloteau, S., Duvert, G., Prato, L., Simon, M., Schuster, K., & Ménard, F. 1996, *A&A*, 309, 493
- Duvert, G., Dutrey, A., Guilloteau, S., Ménard, F., Schuster, K., Prato, L., & Simon, M. 1998, *A&A*, 332, 867
- Greene, T. P., Wilking, B. A., André, P., Young, E. T., & Lada, C. J. 1994, *ApJ*, 434, 614
- Hildebrand, R. H. 1983, *QJRAS*, 24, 267
- Ho, P. T. P., Moran, J. M., & Lo, K.-Y. 2004, *ApJ*, 616, L1

- Jensen, E. L. N., Mathieu, R. D., & Fuller, G. A. 1994, *ApJ*, 429, L29
- 1996, *ApJ*, 458, 312
- Jensen, E. L. N., Koerner, D. W., & Mathieu, R. D. 1996, *AJ*, 111, 2431
- Jensen, E. L. N., & Mathieu, R. D. 1997, *AJ*, 114, 301
- Jensen, E. L. N., & Akeson, R. L. 2003, *ApJ*, 584, 875
- Koerner, D. W., Sargent, A. I., & Beckwith, S. V. W. 1993, *ApJ*, 408, L93
- Lin, D. N. C., & Papaloizou, J. 1993, *Protostars & Planets III*, eds. E. Levy, & M. S. Matthews (Tucson: Univ. Ariz. Press), 749
- Martín, E. L., Montmerle, T., Gregorio-Hetem, J., & Casanova, S. 1998, *MNRAS*, 300, 733
- Mathieu, R. D. 1994, *ARA&A*, 32, 465
- Mathieu, R. D., Adams, F. C., Fuller, G. A., Jensen, E. L. N., Koerner, D. W., & Sargent, A. I. 1995, *AJ*, 109, 2655
- Mathieu, R. D., Martín, E. L., & Magazzù, A. 1996, *BAAS*, 128, 6005
- Mathieu, R. D., Ghez, A. M., Jensen, E. L. N., & Simon, M. 2000, *Protostars and Planets IV*, eds. V. Mannings, A. P. Boss, & S. S. Russell (Tucson: Univ. Ariz. Press), 703
- Nürnberg, D., Brandner, W., Yorke, H. W., & Zinnecker, H. 1998, *A&A*, 330, 549
- Osterloh, M., & Beckwith, S. V. W. 1995, *ApJ*, 439, 288
- Papaloizou, J. C. B., & Pringle, J. E. 1977, *MNRAS*, 181, 441
- Reipurth, B., Chini, R., Krügel, E., Kreysa, E., & Sievers, A. 1993, *A&A*, 273, 221
- Reipurth, B., & Zinnecker, H. 1993, *A&A*, 278, 81
- Scoville, N. Z., Sargent, A. I., Sanders, D. B., Claussen, M. J., Masson, C. R., Lo, K. Y., & Phillips, T. G. 1986, *ApJ*, 303, 416
- Simon, M., et al. 1995, *ApJ*, 443, 625
- Qi, C., Kessler, J. E., Koerner, D. W., Sargent, A. I., & Blake, G. A. 2003, *ApJ*, 597, 986
- Weidenschilling, S. J. 1977, *Ap&SS*, 51, 153
- Weintraub, D. A., Masson, C. R., & Zuckerman, B. 1989, *ApJ*, 344, 915
- Wilner, D. J., & Welch, W. J. 1994, *ApJ*, 427, 898




Fig. 1.— Aperture synthesis images of the millimeter continuum (grayscale) and velocity-integrated CO line emission (contours) in the SR 24 system. The CO contours begin at the 3σ level ($0.82 \text{ Jy km s}^{-1}$) and each step represents a factor of $\sqrt{2}$ in intensity. Negative contours are dashed. The axes are offsets from the position of SR 24 S in arcseconds. The FWHM synthesized beam size is shown in the lower right corner. The stellar positions are marked with crosses of sizes indicating the uncertainty in the absolute HST astrometric positions and the SMA pointing. Gas and dust are present in a circumstellar disk around SR 24 S, while only the gas phase of a circumbinary ring around SR 24 N is detected.




Fig. 2.— The observed SEDs of SR 24 S (circles) and N (triangles). The infrared points are from Greene et al. (1994). The ordinate axis is defined as $\nu L_\nu = 4\pi d^2 \nu F_\nu$. Both SR 24 N and S have typical Class II infrared SEDs, suggesting the presence of inner disks.